### **Sporanox**



**Generic Sporanox** is an antifungal antibiotic. It is used to treat infections caused by fungus, which can invade any part of the body including the lungs, mouth or throat, toenails or fingernails.

Active ingredient: Itraconazole

Brand(s):Canditral

Manufacturer: Glenmark Pharmaceuticals Ltd.

Disease(s):Fungus

#### Sporanox 100mg

|                       | Per pill | Per Pack                         | Order       |  |  |  |
|-----------------------|----------|----------------------------------|-------------|--|--|--|
| 10 tablet             | \$ 7.00  | \$ 70.00                         | Add to cart |  |  |  |
| 20 tablet             | \$ 6.50  | \$ 130.00<br>\$140.00            | Add to cart |  |  |  |
| 30 tablet             | \$ 6.00  | \$ 180.00<br><del>\$210.00</del> | Add to cart |  |  |  |
| Free AirMail shipping |          |                                  |             |  |  |  |
| 40 tablet             | \$ 5.50  | \$ 220.00<br>\$280.00            | Add to cart |  |  |  |
| Free AirMail shipping |          |                                  |             |  |  |  |
| 50 tablet             | \$ 5.00  | \$ 250.00<br><del>\$350.00</del> | Add to cart |  |  |  |
| Free AirMail shipping |          |                                  |             |  |  |  |

## Itraconazole capsule

#### What is this medicine?

ITRACONAZOLE is an antifungal. It is used to treat infections caused by fungus on the nails and in the body.

# What should I tell my health care provider before I take this medicine?

They need to know if you have any of these conditions:

- · heart disease, including angina or heart failure
- · kidney disease or on dialysis
- · liver disease
- lung disease
- an unusual or allergic reaction to itraconazole, or other antifungal medicines, foods, dyes or preservatives
- · pregnant or trying to get pregnant
- · breast-feeding

#### How should I use this medicine?

Take this medicine by mouth with a full glass of water. Take after eating a full meal. If you have a condition called achlorhydria, are taking H2-receptor antagonists or other gastric acid suppressors, you should take this medicine with a cola beverage. Take your doses at regular intervals. Do not take your medicine more often than directed. Do not stop taking except on your doctor's advice.

Talk to your pediatrician regarding the use of this medicine in children. Special care may be needed.

Overdosage: If you think you have taken too much of this medicine