

## Cyclogyl



Generic Cyclogyl is used for dilating the eye before an exam or surgery and preventing the eye from focusing.

Active ingredient: [Cyclopentolate](#)

**Brand(s):**Cyclopent

**Manufacturer:** Sun Pharmaceutical Industries Ltd.

### Cyclogyl 1% 5ml

	Per solution	Per Pack	Order
1 solution	\$ 10.00	<b>\$ 10.00</b>	<a href="#">Add to cart</a>
2 solution	\$ 9.00	<b>\$ 18.00</b> <del>\$20.00</del>	<a href="#">Add to cart</a>
3 solution	\$ 7.00	<b>\$ 21.00</b> <del>\$30.00</del>	<a href="#">Add to cart</a>
4 solution	\$ 6.00	<b>\$ 24.00</b> <del>\$40.00</del>	<a href="#">Add to cart</a>
5 solution	\$ 5.50	<b>\$ 27.50</b> <del>\$50.00</del>	<a href="#">Add to cart</a>
6 solution	\$ 5.00	<b>\$ 30.00</b> <del>\$60.00</del>	<a href="#">Add to cart</a>

## Cyclopentolate 1% solution

### What is this medicine?

CYCLOPENTOLATE is used in the eye to widen your pupils, so they won't respond to light. It is normally used for diagnostic procedures.

### What should my health care professional know before I take this medicine?

They need to know if you have any of these conditions:

- Down's Syndrome
- glaucoma
- wear contact lenses
- an unusual or allergic reaction to cyclopentolate or other medications, foods, dyes or preservatives
- pregnant or trying to become pregnant
- breast-feeding

### How should I use this medicine?

This medicine is only for use in the eye(s). Do not take by mouth. Wash hands before and after use. Tilt the head back slightly and pull the lower eyelid down with the index finger to form a pouch. Squeeze a single drop into the pouch and close your eyes. Apply gentle pressure to the inside corner of your eye for 1 to 3 minutes. If you are using more than one drop, repeat this process with 5 minutes between the drops. Do not touch dropper tip to any surface to prevent contamination. If more than one eye product is being used, the products should be administered at least 5 minutes apart. Use your medicine at regular intervals. Do not use it more often than directed.

Contact your pediatrician or health care professional regarding the use of this medication in children. Special care is needed.

**Overdose:** If you think you have taken too much of this medicine

