

Please click price below and you will be redirected to our pharmacy.

## Kombiglyze XR

5/1000

Metformin and saxagliptin are oral diabetes medicines that help control blood sugar levels. Metformin works by decreasing glucose (sugar) production in the liver and decreasing absorption of glucose by the intestines. Saxagliptin works by regulating the levels of insulin your body produces after eating. Metformin and saxagliptin is a combination medicine that is used together with diet and exercise to improve blood sugar control in adults with type 2 diabetes mellitus. This medicine is not for treating type 1 diabetes.

Active ingredient: [Metformin + Saxagliptin](#)

**Brand(s):**Riix MXR

**Manufacturer:** Dr Reddys

**Disease(s):**Diabetes Type 2

### Kombiglyze XR 1000mg + 5mg

	Per pill	Per Pack	Order
14 tablet	\$ 6.42	<b>\$ 89.95</b>	<a href="#">Add to cart</a>
28 tablet	\$ 5.36	<b>\$ 149.95</b> <del>\$179.90</del>	<a href="#">Add to cart</a>
56 tablet	\$ 4.37	<b>\$ 244.95</b> <del>\$359.80</del>	<a href="#">Add to cart</a>
Free AirMail shipping			
84 tablet	\$ 3.99	<b>\$ 334.95</b> <del>\$539.70</del>	<a href="#">Add to cart</a>
Free AirMail shipping			
112 tablet	\$ 3.62	<b>\$ 404.95</b> <del>\$719.60</del>	<a href="#">Add to cart</a>
Free AirMail shipping			
168 tablet	\$ 3.33	<b>\$ 559.95</b> <del>\$1079.40</del>	<a href="#">Add to cart</a>
Free AirMail shipping			

### What is metformin and saxagliptin?

Metformin and saxagliptin are oral diabetes medicines that help control blood sugar levels. Metformin works by decreasing glucose (sugar) production in the liver and decreasing absorption of glucose by the intestines. Saxagliptin works by regulating the levels of insulin your body produces after eating.

Metformin and saxagliptin is a combination medicine that is used together with diet and exercise to improve blood sugar control in adults with type 2 diabetes mellitus. This medicine is not for treating type 1 diabetes.

Metformin and saxagliptin may also be used for purposes not listed in this medication guide.

### Important Information

You should not use this medicine if you have severe kidney disease or diabetic ketoacidosis (call your doctor for treatment with insulin).

Some people develop lactic acidosis while taking metformin. Early symptoms may get worse over time and this condition can be fatal. **Stop taking this medicine and get emergency medical help if you have even mild symptoms such as:** muscle pain or weakness, trouble breathing, stomach pain, nausea, and feeling very weak or tired.

### Before taking this medicine

