### **Betadine**



Betadine cream is used for treating minor wounds and infections, as well as killing bacteria. Betadine cream is an antiseptic combination. It works by killing sensitive bacteria.

Active ingredient: Providone-iodine

Brand(s):Wokadine

Manufacturer: Wockhardt Ltd. Disease(s):Wound / Infection

#### Betadine 10% 15g

	Per pill	Per Pack	Order
1 tube	\$ 24.95	\$ 24.95	Add to cart
2 tube	\$ 19.98	\$ 39.95 <del>\$49.90</del>	Add to cart
3 tube	\$ 17.65	\$ 52.95 <del>\$74.85</del>	Add to cart
4 tube	\$ 14.99	\$ 59.95 <del>\$99.80</del>	Add to cart
5 tube	\$ 13.99	\$ 69.95 <del>\$124.75</del>	Add to cart

## Povidone-Iodine topical

#### What is this medicine?

Povidone-Iodine cream is used for treating minor wounds and infections, as well as killing bacteria. Povidone-Iodine cream is an antiseptic combination. It works by killing sensitive bacteria.

# What should I tell my health care providers before I take this medicine?

Some medical conditions may interact with Povidone-Iodine cream. Tell your doctor or pharmacist if you have any medical conditions, especially if any of the following apply to you:

- if you are pregnant, planning to become pregnant, or are breastfeeding
- if you are taking any prescription or nonprescription medicine, herbal preparation, or dietary supplement
- · if you have allergies to medicines, foods, or other substances
- if you have serious burns or deep puncture wounds

Some MEDICINES MAY INTERACT with Povidone-Iodine cream. However, no specific interactions are known at this time.

This may not be a complete list of all interactions that may occur. Ask your health care provider if Povidone-Iodine cream may interact with other medicines that you take. Check with your health care provider before you start, stop, or change the dose of any medicine.

#### How should I use this medicine?

Use Povidone-Iodine cream as directed by your doctor. Check the label on the medicine for exact dosing instructions.

- Clean the affected area. Apply a small amount of Povidone-Iodine cream to the affected area. The area may be left uncovered, or you may cover it with a sterile bandage.
- If you miss a dose of Povidone-Iodine cream, use it as soon as