

Elocon



Generic **Elocon** is used to treat the inflammation and itching caused by a number of skin conditions such as allergic reactions, eczema, and psoriasis.

Active ingredient: [Mometasone](#)

Brand(s):Elocon

Manufacturer: ZYG Pharma Pvt. Ltd.

Disease(s):Eczema / Psoriasis / Allergy

Elocon 5g

| | Per tube | Per Pack | Order |
|---------|----------|---------------------------------------|-----------------------------|
| 1 cream | \$ 15.00 | \$ 15.00 | Add to cart |
| 2 cream | \$ 14.00 | \$ 28.00 \$30.00 | Add to cart |
| 3 cream | \$ 13.00 | \$ 39.00 \$45.00 | Add to cart |
| 4 cream | \$ 12.00 | \$ 48.00 \$60.00 | Add to cart |
| 5 cream | \$ 11.00 | \$ 55.00 \$75.00 | Add to cart |
| 6 cream | \$ 10.00 | \$ 60.00 \$90.00 | Add to cart |

Mometasone 0.1% cream

What is this medicine?

MOMETASONE is a corticosteroid. It is used to treat skin problems that may cause itching, redness, and swelling.

What should my health care professional know before I take this medicine?

They need to know if you have any of these conditions:

- acne or rosacea
- any type of active infection
- large areas of burned or damaged skin
- skin wasting or thinning
- an unusual or allergic reaction to mometasone, steroids, other medicines, foods, dyes, or preservatives
- pregnant or trying to get pregnant
- breast-feeding

How should I use this medicine?

This medicine is for external use only. Do not take by mouth. Wash your hands before and after use. Apply a thin film to the affected area and rub in gently. Do not bandage or wrap the skin being treated unless directed to do so by your doctor or health care professional. Do not use on healthy skin or over large areas of skin. Do not get this medicine in your eyes. If you do, rinse it out with plenty of cool tap water.

Use your doses at regular intervals. Do not use your medicine more often than directed or for a longer period of time than ordered by your doctor or health care professional. To do so may increase the chance of side effects.

Talk to your pediatrician regarding the use of this medicine in children

