

## Diclegis



**Diclegis** is a delayed-release tablet containing a combination of doxylamine and pyridoxine. Doxylamine is an antihistamine that reduces the effects of natural chemical histamine in the body. Pyridoxine is a form of vitamin B (B6). **Diclegis** is used to treat nausea and vomiting during pregnancy, sometimes called morning sickness.

Active ingredient: [Pyridoxine And Doxylamine](#)

**Brand(s):** Pregnidoxin Nu Tab

**Manufacturer:** Ucb

**Disease(s):** Nausea / Vomiting / Vomiting During Pregnancy / Morning Sickness

### Diclegis 10mg + 10mg

	Per pill	Per Pack	Order
30 tablet	\$ 2.67	<b>\$ 79.95</b>	<a href="#">Add to cart</a>
60 tablet	\$ 2.33	<b>\$ 139.95</b> <del>\$159.90</del>	<a href="#">Add to cart</a>
90 tablet	\$ 2.22	<b>\$ 199.95</b> <del>\$239.85</del>	<a href="#">Add to cart</a>
Free AirMail shipping			
120 tablet	\$ 2.08	<b>\$ 249.95</b> <del>\$319.80</del>	<a href="#">Add to cart</a>
Free AirMail shipping			
180 tablet	\$ 2.00	<b>\$ 359.95</b> <del>\$479.70</del>	<a href="#">Add to cart</a>
Free AirMail shipping			

## Doxylamine and pyridoxine tablets

### What is this medicine?

DOXYLAMINE is an antihistamine that reduces the effects of natural chemical histamine in the body.

PYRIDOXINE is a form of vitamin B (B6).

Doxylamine and pyridoxine is a combination medicine used to treat nausea and vomiting during pregnancy, sometimes called morning sickness.

### What should I tell my health care providers before I take this medicine?

Do not use doxylamine and pyridoxine if you have used an MAO inhibitor in the past 14 days. A dangerous drug interaction could occur. MAO inhibitors include isocarboxazid, linezolid, methylene blue injection, phenelzine, rasagiline, selegiline, tranylcypromine, and others.

You should not use this medicine if you are allergic to doxylamine or pyridoxine, or to other antihistamines such as Benadryl or Dramamine.

To make sure this medicine is safe for you, tell your doctor if you have:

- asthma or other breathing disorder;
- glaucoma, increased pressure inside your eye;
- blockage in your digestive tract (stomach or intestines);
- a stomach ulcer; or
- bladder obstruction or other urination problems.

